

Synaptotagmin4

Cat.No. 105 043; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

| | |
|----------------------------|---|
| Reconstitution/ Storage | 50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin was added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet. |
| Applications | WB: 1 : 100 up to 1 : 1000 (AP staining) IP: yes ICC: not recommended (see remarks) IHC: not recommended IHC-P: not tested yet |
| Immunogen | Recombinant protein corresponding to AA 40 to 151 from rat Synaptotagmin4 (UniProt Id: P50232) |
| Reactivity | Reacts with: rat (P50232), mouse (P40749). Other species not tested yet. |
| Specificity | K.O. validated |
| Matching control | 105-4P |
| Remarks | ICC: Cat. no. 105 143 is recommended. |

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Up to now at least 17 synaptotagmins have been identified. **Synaptotagmin4** is composed of a vesicular, a transmembrane and two C2 domains. Only the C2B domain is able to bind calcium. In the C2A domain one of the calcium binding aspartates has been substituted for serine leading to a loss of its binding capabilities.

The localization of synaptotagmin4 is still under discussion. A localization to synaptic vesicles (SVs) has been postulated but more recent studies suggest that it is present in the Golgi compartment, in distal parts of neurites and on large dense core vesicles (LDCVs) of NGF differentiated PC12 cells.

Selected References for 105 043

Involvement of complexin 2 in docking, locking and unlocking of different SNARE complexes during sperm capacitation and induced acrosomal exocytosis.

Tsai PS, Brewis IA, van Maaren J, Gadella BM
PloS one (2012) 73: e32603. . **WB, ICC; tested species: pig**

Cell therapy modulates expression of Tax1-binding protein 1 and synaptotagmin IV in a model of optic nerve lesion.
Mesentier-Louro LA, Coronel J, Zaverucha-do-Valle C, Mencia A, Paredes BD, Abdelhay E, Mendez-Otero R, Santiago MF
Investigative ophthalmology & visual science (2012) 538: 4720-9. . **WB, IHC**

Loss of synaptotagmin IV results in a reduction in synaptic vesicles and a distortion of the Golgi structure in cultured hippocampal neurons.

Arthur CP, Dean C, Pagratis M, Chapman ER, Stowell MH
Neuroscience (2010) 1671: 135-42. . **WB, IP**

Synaptotagmin-IV modulates synaptic function and long-term potentiation by regulating BDNF release.

Dean C, Liu H, Dunning FM, Chang PY, Jackson MB, Chapman ER
Nature neuroscience (2009) 126: 767-76. . **WB, ICC; IP**

Synaptotagmin IV: a multifunctional regulator of peptidergic nerve terminals.

Zhang Z, Bhalla A, Dean C, Chapman ER, Jackson MB
Nature neuroscience (2009) 122: 163-71. . **WB, EM**

Synaptotagmin 4 Regulates Pancreatic β Cell Maturation by Modulating the Ca²⁺ Sensitivity of Insulin Secretion Vesicles.

Huang C, Walker EM, Dadi PK, Hu R, Xu Y, Zhang W, Sanavia T, Mun J, Liu J, Nair GG, Tan HYA, et al.
Developmental cell (2018) 453: 347-361.e5. . **WB, IHC; tested species: mouse**

Distinct subsets of Syt-IV/BDNF vesicles are sorted to axons versus dendrites and recruited to synapses by activity.

Dean C, Liu H, Staudt T, Stahlberg MA, Vingill S, Bückers J, Kamin D, Engelhardt J, Jackson MB, Hell SW, Chapman ER, et al.
The Journal of neuroscience : the official journal of the Society for Neuroscience (2012) 3216: 5398-413. . **WB, ICC**

REST/NRSF drives homeostatic plasticity of inhibitory synapses in a target-dependent fashion.

Prestigio C, Ferrante D, Marte A, Romei A, Lignani G, Onofri F, Valente P, Benfenati F, Baldelli P
eLife (2021) 10: . . **WB; tested species: mouse**

Synaptotagmin oligomerization is essential for calcium control of regulated exocytosis.

Bello OD, Jouannot O, Chaudhuri A, Stroeve E, Coleman J, Volynski KE, Rothman JE, Krishnakumar SS
Proceedings of the National Academy of Sciences of the United States of America (2018) 11532: E7624-E7631. . **WB; tested species: rat**

Reduced insulin secretion correlates with decreased expression of exocytotic genes in pancreatic islets from patients with type 2 diabetes.

Andersson SA, Olsson AH, Esguerra JL, Heimann E, Ladenvall C, Edlund A, Salehi A, Taneera J, Degerman E, Groop L, Ling C, et al.
Molecular and cellular endocrinology (2012) 3641-2: 36-45. . **WB**

REST/NRSF governs the expression of dense-core vesicle gliosecretion in astrocytes.

Prada I, Marchaland J, Podini P, Magrassi L, D'Alessandro R, Bezzi P, Meldolesi J
The Journal of cell biology (2011) 1933: 537-49. . **WB**

Access the online factsheet including applicable protocols
at <https://sysy.com/product/105043> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.