

Synaptotagmin1 luminal domain

Cat.No. 105 221; Monoclonal mouse antibody, 200 µl hybridoma supernatant (lyophilized)

Data Sheet

Reconstitution/ Storage	200 µl hybridoma supernatant, lyophilized. For reconstitution add 200 µl H ₂ O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 up to 1 : 10000 (AP staining) (see remarks) IP: yes ICC: 1 : 100 (see remarks) IHC: 1 : 500 (see remarks) IHC-P: not tested yet
Clone	604.1
Subtype	IgG3 (κ light chain)
Immunogen	Synthetic peptide corresponding to residues near the amino terminus of rat Synaptotagmin1 (UniProt Id: P21707)
Reactivity	Reacts with: rat (P21707), mouse (P46096). No signal: zebrafish. Other species not tested yet.
Specificity	K.O. validated PubMed: 26195798
Remarks	WB: Only detects rat Synaptotagmin1 in westernblots. ICC: Antibody is suitable for the labeling of recycling vesicles of living rat and mouse neurons. It detects PFA fixed rat Synaptotagmin1, but is negative on PFA fixed mouse Synaptotagmin1. IHC: Antibody detects PFA fixed rat Synaptotagmin1, but is negative on PFA fixed mouse Synaptotagmin1.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Synaptotagmin1, also known as **p65**, is an integral membrane glycoprotein of neuronal synaptic vesicles and secretory granules of neuroendocrine cells that is widely (but not ubiquitously) expressed in the central and peripheral nervous system. It has a variable N-terminal domain that is exposed to the lumen of the vesicle and a conserved cytoplasmic tail that contains two Ca²⁺-binding C2-domains. Ca²⁺-binding to synaptotagmin triggers exocytosis of synaptic vesicles, thus linking Ca²⁺-influx during depolarization to neurotransmitter release.

Luminal antibodies were used in living neurons to label synaptic vesicles from the outside via endocytotic uptake.

Selected References for 105 221

Structural elements that underlie Doc2β function during asynchronous synaptic transmission.

Xue R, Gaffaney JD, Chapman ER

Proceedings of the National Academy of Sciences of the United States of America (2015) 11231: E4316-25. . **WB, ICC; KO verified; tested species: mouse**

Storage and uptake of D-serine into astrocytic synaptic-like vesicles specify gliotransmission.

Martineau M, Shi T, Puyal J, Knolhoff AM, Dulong J, Gasnier B, Klingauf J, Sweedler JV, Jahn R, Mothet JP

The Journal of neuroscience : the official journal of the Society for Neuroscience (2013) 338: 3413-23. . **IP, IHC; tested species: rat**

SV2B regulates synaptotagmin 1 by direct interaction.

Lazzell DR, Belizaire R, Thakur P, Sherry DM, Janz R

The Journal of biological chemistry (2004) 27950: 52124-31. . **IP, WB; tested species: mouse**

STED microscopy reveals that synaptotagmin remains clustered after synaptic vesicle exocytosis.

Willig KI, Rizzoli SO, Westphal V, Jahn R, Hell SW

Nature (2006) 4407086: 935-9. . **UPTAKE**

Disease-linked mutations in Munc18-1 deplete synaptic Doc2.

Guiberson NGL, Black LS, Haller JE, Brukner A, Abramov D, Ahmad S, Xie YX, Sharma M, Burré J

Brain : a journal of neurology (2024) : . . **UPTAKE; tested species: mouse**

Targeted stabilization of Munc18-1 function via pharmacological chaperones.

Abramov D, Guiberson NGL, Daab A, Na Y, Petsko GA, Sharma M, Burré J

EMBO molecular medicine (2020) : e12354. . **UPTAKE; tested species: mouse**

Mechanism-based rescue of Munc18-1 dysfunction in varied encephalopathies by chemical chaperones.

Guiberson NGL, Pineda A, Abramov D, Kharel P, Carnazza KE, Wragg RT, Dittman JS, Burré J

Nature communications (2018) 91: 3986. . **UPTAKE; tested species: mouse**

Loss of Doc2-Dependent Spontaneous Neurotransmission Augments Glutamatergic Synaptic Strength.

Ramirez DMO, Crawford DC, Chanaday NL, Trauterman B, Monteggia LM, Kavalali ET

The Journal of neuroscience : the official journal of the Society for Neuroscience (2017) 3726: 6224-6230. . **UPTAKE; tested species: rat**

BDNF enhances spontaneous and activity-dependent neurotransmitter release at excitatory terminals but not at inhibitory terminals in hippocampal neurons.

Shinoda Y, Ahmed S, Ramachandran B, Bharat V, Brockelt D, Altas B, Dean C

Frontiers in synaptic neuroscience (2014) 6: 27. . **ICC; tested species: rat**

CSPα knockout causes neurodegeneration by impairing SNAP-25 function.

Sharma M, Burré J, Bronk P, Zhang Y, Xu W, Südhof TC

The EMBO journal (2012) 314: 829-41. . **WB; tested species: mouse**

Access the online factsheet including applicable protocols at <https://sysy.com/product/105221> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.