

Syntaxin6

Cat.No. 110 062; Polyclonal rabbit antibody, 200 µl antiserum (lyophilized)

Data Sheet

Reconstitution/ Storage	200 µl antiserum, lyophilized. For reconstitution add 200 µl H ₂ O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 up to 1 : 10000 (AP staining) IP: yes ICC: 1 : 500 IHC: not tested yet IHC-P: not tested yet
Immunogen	Recombinant protein corresponding to AA 1 to 235 from human Syntaxin6 (UniProt Id: O43752)
Reactivity	Reacts with: human (O43752), rat (Q63635), mouse (Q9JJK1), hamster, chicken. Other species not tested yet.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Syntaxin 6, a member of the SNARE family of proteins, localizes to the trans-Golgi network and endosomal structures. It is widely expressed with relatively higher expression in brain, lung and kidney.

In contrast to other SNARE proteins it has a broad variety of interaction partners depending on the cell type examined. It is therefore suggested that syntaxin 6 is involved in multiple membrane-trafficking events, e.g. the transport from the early endosome to the trans-Golgi network.

Syntaxin 6 has been identified as a Qc SNARE due to its close homology to the C-terminal SNARE domain of SNAP 25.

Selected References for 110 062

A novel syntaxin 6-interacting protein, SHIP164, regulates syntaxin 6-dependent sorting from early endosomes.
Otto GP, Razi M, Morvan J, Stenner F, Tooze SA
Traffic (Copenhagen, Denmark) (2010) 115: 688-705. . **IP, WB, ICC; tested species: human**

A trap mutant reveals the physiological client spectrum of TRC40.
Coy-Vergara J, Rivera-Monroy J, Urlaub H, Lenz C, Schwappach B
Journal of cell science (2019) 13213: . . **WB, ICC; tested species: human**

SNAREs define targeting specificity of trafficking vesicles by combinatorial interaction with tethering factors.
Koike S, Jahn R
Nature communications (2019) 101: 1608. . **WB, ICC; tested species: human**

Probing and manipulating intracellular membrane traffic by microinjection of artificial vesicles.
Koike S, Jahn R
Proceedings of the National Academy of Sciences of the United States of America (2017) 11446: E9883-E9892. . **WB, ICC; tested species: human**

Dynamic GLUT4 sorting through a syntaxin-6 compartment in muscle cells is derailed by insulin resistance-causing ceramide.
Foley KP, Klip A
Biology open (2014) 35: 314-25. . **WB, ICC**

Dual roles of the mammalian GARP complex in tethering and SNARE complex assembly at the trans-golgi network.
Pérez-Victoria FJ, Bonifacino JS
Molecular and cellular biology (2009) 2919: 5251-63. . **WB, ICC**

Trans-Golgi network syntaxin 10 functions distinctly from syntaxins 6 and 16.
Wang Y, Tai G, Lu L, Johannes L, Hong W, Tang BL
Molecular membrane biology () 224: 313-25. . **WB, ICC**

Proteomic analysis reveals the composition of glutamatergic organelles of auditory inner hair cell.
Cepeda AP, Ninov M, Neef J, Parfentev I, Kusch K, Reisinger E, Jahn R, Moser T, Urlaub H
Molecular & cellular proteomics : MCP (2023) : 100704. . **IHC; tested species: mouse**

Endoplasmic reticulum stress impedes regulated secretion by governing key exocytotic and granulogenic molecular switches.
Mukherjee M, Mukherjee C, Ghosh V, Jain A, Sadhukhan S, Dagar S, Sahu BS
Journal of cell science (2024) 1376: . . **WB; tested species: rat**

VPS13B is localized at the interface between Golgi cisternae and is a functional partner of FAM177A1.
Ugur B, Schueder F, Shin J, Hanna MG, Wu Y, Leonzino M, Su M, McAdow AR, Wilson C, Postlethwait J, Solnica-Krezel L, et al.
The Journal of cell biology (2024) 22312: . . **WB; tested species: human**

VPS13B is localized at the cis-trans Golgi complex interface and is a functional partner of FAM177A1.
Ugur B, Schueder F, Shin J, Hanna MG, Wu Y, Leonzino M, Su M, McAdow AR, Wilson C, Postlethwait J, Solnica-Krezel L, et al.
bioRxiv : the preprint server for biology (2023) : . . **WB; tested species: zebrafish**

Access the online factsheet including applicable protocols
at <https://sysy.com/product/110062> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.