

## SV2 B

Cat.No. 119 111; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

### Data Sheet

Reconstitution/ Storage	100 µg purified IgG, lyophilized. Albumin and azide were added for stabilization. For <b>reconstitution</b> add 100 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 1000 (AP staining) (see remarks) <b>IP:</b> yes <b>ICC:</b> 1 : 2000 up to 1 : 5000 <b>IHC:</b> 1 : 200 up to 1 : 500 <b>IHC-P:</b> 1 : 500 <b>DNA-PAINT:</b> yes (see remarks)
Clone	246E8
Subtype	IgG2b (κ light chain)
Immunogen	Synthetic peptide corresponding to AA 2 to 17 from rat SV2B (UniProt Id: Q63564)
Reactivity	Reacts with: human (Q7L1I2), rat (Q63564), mouse (Q8BG39). Other species not tested yet.
Matching control	119-1P
Remarks	<b>WB:</b> SV2 B aggregates after boiling, making it necessary to run SDS-PAGE with non-boiled samples. <b>DNA-PAINT:</b> This antibody has been successfully used for DNA-PAINT application (see Unterauer et al., 2024; <a href="https://pubmed.ncbi.nlm.nih.gov/38552614/">PMID: 38552614</a> ).

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

## Background

**SV2s** (synaptic vesicle protein 2) are integral membrane glycoproteins present in all synaptic vesicles. They have 12 transmembrane domains predicted by sequence analysis. There are three characterized isoforms, SV2 A, SV2 B and SV2 C. SV2 A is expressed ubiquitously throughout the brain and is probably involved in the maintenance of a pool of synaptic vesicles competent for calcium-stimulated exocytosis. **SV2 B** has a more restricted distribution with varying degrees of coexpression with SV2 A. SV2 C is more closely related to SV2 A but shows a very restricted expression pattern. The highest expression levels were observed in phylogenetically old brain areas like pallidum, the midbrain and the olfactory bulb. SV2 expression has also been observed in other organs. In kidney it localizes to podocytes.

## Selected References for 119 111

Loss of photoreceptors results in upregulation of synaptic proteins in bipolar cells and amacrine cells.

Dagar S, Nagar S, Goel M, Cherukuri P, Dhingra NK  
PloS one (2014) 93: e90250. . **IHC, WB; tested species: mouse**

Spatial proteomics in neurons at single-protein resolution.  
Unterauer EM, Shetab Boushehri S, Jevdokimenko K, Masullo LA, Ganji M, Sograte-Idrissi S, Kowalewski R, Strauss S, Reinhardt SCM, Perovic A, Marr C, et al.  
Cell (2024) 1877: 1785-1800.e16. . **DNA\_PAINT; tested species: rat**

Proteomic analysis reveals the composition of glutamatergic organelles of auditory inner hair cell.  
Cepeda AP, Ninov M, Neef J, Parfentev I, Kusch K, Reisinger E, Jahn R, Moser T, Urlaub H  
Molecular & cellular proteomics : MCP (2023) : 100704. . **IHC; tested species: mouse**

Tuning of glutamate, but not GABA, release by an intra-synaptic vesicles APP domain whose function can be modulated by β- or α-secretase cleavage.  
Yao W, Tambini MD, Liu X, D'Adamo L  
The Journal of neuroscience : the official journal of the Society for Neuroscience (2019) : . . **WB; tested species: mouse**

Botulinum Neurotoxins A, B, C, E, and F preferentially enter cultured human motor neurons compared to other cultured human neuronal populations.  
Pellett S, Tepp WH, Johnson EA  
FEBS letters (2019) : . . **WB; tested species: human**

Ubiquitin-Synaptobrevin Fusion Protein Causes Degeneration of Presynaptic Motor Terminals in Mice.  
Liu Y, Li H, Sugiura Y, Han W, Gallardo G, Khvotchev M, Zhang Y, Kavalali ET, Südhof TC, Lin W  
The Journal of neuroscience : the official journal of the Society for Neuroscience (2015) 3533: 11514-31. . **WB**

## Selected General References

SV2 modulates the size of the readily releasable pool of secretory vesicles.

Xu T, Bajjalieh SM  
Nature cell biology (2001) 38: 691-8. .

Genetics of synaptic vesicle function: toward the complete functional anatomy of an organelle.  
Fernández-Chacón R, Südhof TC  
Annual review of physiology (1999) 61: 753-76. .

The synaptic vesicle cycle: a cascade of protein-protein interactions.  
Südhof TC  
Nature (1995) 3756533: 645-53. .

Synaptic vesicles and exocytosis.  
Jahn R, Südhof TC  
Annual review of neuroscience (1994) 17: 219-46. .

Access the online factsheet including applicable protocols  
at <https://sysy.com/product/119111> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C.

### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.