

α/β Synuclein

Cat.No. 128 002; Polyclonal rabbit antibody, 200 µl antiserum (lyophilized)

Data Sheet

Reconstitution/ Storage	200 µl antiserum, lyophilized. For reconstitution add 200 µl H ₂ O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP staining) IP: yes ICC: 1 : 500 IHC: 1 : 500 IHC-P: 1 : 1000 up to 1 : 2000
Immunogen	Synthetic peptide corresponding to AA 2 to 25 from human α-Synuclein (UniProt Id: P37840)
Reactivity	Reacts with: rat (P37377, Q63754), mouse (O55042, Q91ZZ3), zebrafish, human (P37840, Q16143). Other species not tested yet.
Specificity	Recognizes α synuclein and β synuclein.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Synuclein proteins are produced by three genes. They share structural resemblance to apolipoproteins, but are abundant in the neuronal cytosol and present in enriched amounts at presynaptic terminals.

Synucleins have been specifically implicated in three diseases: Alzheimer's (AD), Parkinson's (PD) and breast cancer. In AD, a peptide derived from α-synuclein forms an intrinsic component of plaque amyloid. In PD, an α-synuclein allele is genetically linked to several independent familial cases, and the protein appears to accumulate in Lewy bodies. In breast cancer, increased expression of γ-synuclein correlates with disease progression.

In songbirds, α-synuclein expression is correlated with plasticity in the developing song control system. Although the normal function of synucleins is unknown, a role in synaptic plasticity seems likely.

Selected References for 128 002

Mitochondrial translocation of alpha-synuclein is promoted by intracellular acidification.

Cole NB, Dieuliis D, Leo P, Mitchell DC, Nussbaum RL

Experimental cell research (2008) 31410: 2076-89. . **WB, ICC, EM**

Composition of isolated synaptic boutons reveals the amounts of vesicle trafficking proteins.

Wilhelm BG, Mandad S, Truckenbrodt S, Kröhnert K, Schäfer C, Rammner B, Koo SJ, Claßen GA, Krauss M, Haucke V, Urlaub H, et al.

Science (New York, N.Y.) (2014) 3446187: 1023-8. . **WB, ICC, IHC; tested species: mouse, rat**

Spatial proteomics in neurons at single-protein resolution.

Unterauer EM, Shetab Boushehri S, Jevdokimenko K, Masullo LA, Ganji M, Sograte-Idrissi S, Kowalewski R, Strauss S, Reinhardt SCM, Perovic A, Marr C, et al.

Cell (2024) 1877: 1785-1800.e16. . **DNA_PAINT; tested species: rat**

One-step nanoscale expansion microscopy reveals individual protein shapes.

Shaib AH, Chouaib AA, Chowdhury R, Altendorf J, Mihaylov D, Zhang C, Krah D, Imani V, Spencer R KW, Georgiev SV, Mougios N, et al.

Nature biotechnology (2024) : . . **EXM; tested species: rat**

Trafficking proteins show limited differences in mobility across different postsynaptic spines.

Mougios N, Opazo F, Rizzoli SO, Reshetniak S

iScience (2023) 262: 105971. . **ICC; tested species: rat**

DOPAL initiates αSynuclein-dependent impaired proteostasis and degeneration of neuronal projections in Parkinson's disease.

Masato A, Plotegher N, Terrin F, Sandre M, Faustini G, Thor A, Adams S, Berti G, Cogo S, De Lazzari F, Fontana CM, et al.

NPJ Parkinson's disease (2023) 91: 42. . **WB; tested species: mouse, rat**

A nanobody-based fluorescent reporter reveals human α-synuclein in the cell cytosol.

Gerdes C, Waal N, Offner T, Fornasiero EF, Wender N, Verbarq H, Manzini I, Trenkwalder C, Mollenhauer B, Strohäker T, Zweckstetter M, et al.

Nature communications (2020) 111: 2729. . **ICC; tested species: rat**

Over-expression of alpha-synuclein in the nervous system enhances axonal degeneration after peripheral nerve lesion in a transgenic mouse strain.

Siebert H, Kahle PJ, Kramer ML, Isik T, Schlüter OM, Schulz-Schaeffer WJ, Brück W

Journal of neurochemistry (2010) 1144: 1007-18. . **WB**

Brain alpha-synuclein accumulation in multiple system atrophy, Parkinson's disease and progressive supranuclear palsy: a comparative investigation.

Tong J, Wong H, Guttman M, Ang LC, Forno LS, Shimadzu M, Rajput AH, Muentner MD, Kish SJ, Hornykiewicz O, Furukawa Y, et al.

Brain : a journal of neurology (2010) 133Pt 1: 172-88. . **WB; tested species: human**

Access the online factsheet including applicable protocols
at <https://sysy.com/product/128002> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.