

RIM1

Cat.No. 140 013; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin and azide were added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: yes limited (see remarks) IP: not tested yet ICC: 1 : 500 up to 1 : 1000 IHC: 1 : 500 IHC-P: not tested yet
Immunogen	Recombinant protein corresponding to AA 207 to 366 from rat Rim1 (UniProt Id: Q9JIR4)
Reactivity	Reacts with: rat (Q9JIR4), mouse (Q99NE5). Other species not tested yet.
Specificity	K.D. validated PubMed: 29230050
Remarks	WB: Cat. no. 140 023 is recommended for westernblotting.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

RIMs are presynaptic active zone proteins that regulate Ca²⁺ triggered release of neurotransmitters. RIM 1α and RIM 2α are composed of an N-terminal zinc-finger domain, a central PDZ domain and two C-terminal C2 domains that are separated by long alternatively spliced sequences. RIM 1α is a putative Rab 3a effector and has been shown to interact with other active zone proteins like Munc13-1, ERC 1b, ERC 2 and α-liprins. Deletion of RIM 1α in mice impaired neurotransmitter release without changing the structure of the synapse.

Selected References for 140 013

cAMP-EPAC-PKCε-RIM1α signaling regulates presynaptic long-term potentiation and motor learning. Wang XT, Zhou L, Dong BB, Xu FX, Wang DJ, Shen EW, Cai XY, Wang Y, Wang N, Ji SJ, Chen W, et al. *eLife* (2023) 12: . . **IP, ICC; tested species: mouse**

Analysis of RIM Expression and Function at Mouse Photoreceptor Ribbon Synapses. Löhner M, Babai N, Müller T, Gierke K, Atorf J, Joachimsthaler A, Peukert A, Martens H, Feigenspan A, Kremers J, Schoch S, et al. *The Journal of neuroscience : the official journal of the Society for Neuroscience* (2017) 3733: 7848-7863. . **WB, IHC; tested species: mouse**

Postsynaptic RIM1 modulates synaptic function by facilitating membrane delivery of recycling NMDARs in hippocampal neurons.

Wang J, Lv X, Wu Y, Xu T, Jiao M, Yang R, Li X, Chen M, Yan Y, Chen C, Dong W, et al. *Nature communications* (2018) 91: 2267. . **IHC; tested species: mouse**

Synaptic weight set by Munc13-1 supramolecular assemblies.

Sakamoto H, Ariyoshi T, Kimpara N, Sugao K, Taiko I, Takikawa K, Asanuma D, Namiki S, Hirose K. *Nature neuroscience* (2018) 211: 41-49. . **ICC; KD verified; tested species: rat**

Regulation of density of functional presynaptic terminals by local energy supply.

Zhou H, Liu G

Molecular brain (2015) 8: 42. . **ICC; tested species: rat**

Molecular in situ topology of Aczonin/Piccolo and associated proteins at the mammalian neurotransmitter release site.

Limbach C, Laue MM, Wang X, Hu B, Thiede N, Hultqvist G, Kilimann MW

Proceedings of the National Academy of Sciences of the United States of America (2011) 10831: E392-401. . **WB**

Selected General References

Genomic definition of RIM proteins: evolutionary amplification of a family of synaptic regulatory proteins.

Wang Y, Südhof TC

Genomics (2003) 812: 126-37. .

RIM1alpha is required for presynaptic long-term potentiation.

Castillo PE, Schoch S, Schmitz F, Südhof TC, Malenka RC

Nature (2002) 4156869: 327-30. .

RIM1alpha forms a protein scaffold for regulating neurotransmitter release at the active zone.

Schoch S, Castillo PE, Jo T, Mukherjee K, Geppert M, Wang Y, Schmitz F, Malenka RC, Südhof TC

Nature (2002) 4156869: 321-6. .

The RIM/NIM family of neuronal C2 domain proteins. Interactions with Rab3 and a new class of Src homology 3 domain proteins.

Wang Y, Sugita S, Südhof TC

The Journal of biological chemistry (2000) 27526: 20033-44. .

Rim is a putative Rab3 effector in regulating synaptic-vesicle fusion.

Wang Y, Okamoto M, Schmitz F, Hofmann K, Südhof TC

Nature (1997) 3886642: 593-8. .

Access the online factsheet including applicable protocols at <https://sysy.com/product/140013> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.