

## GOSR1

Cat.No. 179-0P; control protein, 100 µg protein (lyophilized)

### Data Sheet

Reconstitution/ Storage	100 µg protein, lyophilized. For <b>reconstitution</b> add 100 µl H <sub>2</sub> O to get a 1mg/ml solution in TBS. Then aliquot and store at -20°C to -80°C until use. Control proteins should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Immunogen	Recombinant protein corresponding to AA 2 to 155 from rat GOSR1 (UniProt Id: Q62931)
Recommended dilution	Optimal concentrations should be determined by the end-user.
Matching antibodies	179 002, 179 003
Remarks	This control protein consists of the recombinant protein (aa 2 - 155 of rat GOSR 1) that has been used for immunization. It has been tested in preadsorption experiments and blocks efficiently and specifically the corresponding signal in Western blots. The amount of protein needed for efficient blocking depends on the titer and on the affinity of the antibody to the antigen.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

### Background

**GOSR 1** also known as **GOS 28** and **GS 28** is a 28 kDa membrane protein on the surface of the Golgi compartment. It has been shown to serve as a vSNARE in ER to Golgi transport. In co-immunoprecipitation studies GOSR 1 was identified as a member of a SNARE complex consisting of syntaxin 5, membrin (GS 27 GOSR 2, GOS 27), rbet1 and rsec22. A more detailed analysis revealed two subcomplexes within this complex. One contains syntaxin 5 (mainly the shorter 35 kDa variant) and GOSR 1 whereas the other is composed of syntaxin 5 (35 and 42 kDa variant), membrin, rsec22 and rbet1. It has been suggested that these complexes mediate the fusion of ER-derived vesicles with vesicular tubular clusters (VTC), and the fusion of VTCs to form the cis-Golgi compartment.

### Selected General References

- Targeting of Arf-1 to the early Golgi by membrin, an ER-Golgi SNARE.  
Honda A, Al-Awar OS, Hay JC, Donaldson JG  
The Journal of cell biology (2005) 1687: 1039-51. .
- Counterstream distribution of two distinct SNARE complexes mediating transport within the Golgi stack.  
Volchuk A, Ravazzola M, Perrelet A, Eng WS, Di Liberto M, Varlamov O, Fukasawa M, Engel T, Söllner TH, Rothman JE, Orci L, et al.  
Molecular biology of the cell (2004) 154: 1506-18. .
- The SNARE motif contributes to rbet1 intracellular targeting and dynamics independently of SNARE interactions.  
Joglekar AP, Xu D, Rigotti DJ, Fairman R, Hay JC  
The Journal of biological chemistry (2003) 27816: 14121-33. .
- Sequential tethering of Golgins and catalysis of SNAREpin assembly by the vesicle-tethering protein p115.  
Shorter J, Beard MB, Seemann J, Dirac-Svejstrup AB, Warren G  
The Journal of cell biology (2002) 1571: 45-62. .
- Subunit structure of a mammalian ER/Golgi SNARE complex.  
Xu D, Joglekar AP, Williams AL, Hay JC  
The Journal of biological chemistry (2000) 27550: 39631-9. .
- SNARE membrane trafficking dynamics in vivo.  
Chao DS, Hay JC, Winnick S, Prekeris R, Klumperman J, Scheller RH  
The Journal of cell biology (1999) 1445: 869-81. .
- Localization, dynamics, and protein interactions reveal distinct roles for ER and Golgi SNAREs.  
Hay JC, Klumperman J, Oorschot V, Steegmaier M, Kuo CS, Scheller RH  
The Journal of cell biology (1998) 1417: 1489-502. .
- Role of vesicle-associated syntaxin 5 in the assembly of pre-Golgi intermediates.  
Rowe T, Dascher C, Bannykh S, Plutner H, Balch WE  
Science (New York, N.Y.) (1998) 2795351: 696-700. .

Access the online factsheet including applicable protocols at <https://sysy.com/product/179-0P> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C.

### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.