

Parvalbumin

Cat.No. 195 308; Recombinant Guinea pig antibody, 50 µg recombinant IgG (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg purified recombinant IgG, lyophilized. Albumin and azide were added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 500 (AP staining) (see remarks) IP: not tested yet ICC: 1 : 500 IHC: 1 : 1000 up to 1 : 2000 IHC-P: 1 : 1000 up to 1 : 4000
Clone	Gp58E1
Subtype	IgG2 (κ light chain)
Immunogen	Full-length recombinant rat Parvalbumin (UniProt Id: P02625)
Reactivity	Reacts with: rat (P02625), mouse (P32848). No signal: zebrafish. Other species not tested yet.
Remarks	This antibody is a chimeric antibody based on the monoclonal mouse antibody clone 58E1. The constant regions of the heavy and light chains have been replaced by guinea pig specific sequences. Therefore, the antibody can be used with standard anti-guinea pig secondary reagents. The antibody has been expressed in mammalian cells. WB: Due to the small size of the protein, a tricine gel is recommended. The rabbit polyclonal antiserum (cat. no. 195 002) is more sensitive and recommended for westernblotting.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Parvalbumin is a small, acidic calcium binding protein and belongs to the family of EF hand proteins. The protein is found in skeletal muscle and the brain of vertebrates where it locates to a specific population of GABAergic interneurons. This subset of neurons may contribute to maintaining the balance between excitation and inhibition in the cortex and the hippocampus.

Selected References for 195 308

Distinct ventral hippocampal inhibitory microcircuits regulating anxiety and fear behaviors.
Li K, Koukoutselos K, Sakaguchi M, Ciochi S
Nature communications (2024) 151: 8228. . **IHC; tested species: mouse**

Pleiotrophin Overexpression Reduces Adolescent Ethanol Consumption and Modulates Ethanol-Induced Glial Responses and Changes in the Perineuronal Nets in the Mouse Hippocampus.
Galán-Llario M, Rodríguez-Zapata M, Fontán-Baselga T, Cañeque-Rufo H, García-Guerra A, Fernández B, Gramage E, Herradón G
CNS neuroscience & therapeutics (2024) 3012: e70159. . **IHC; tested species: mouse**

Selected General References

Quantitative analysis of parvalbumin-immunoreactive cells in the human epileptic hippocampus.
Andrioli A et al. Neuroscience (2007) PubMed:17850980

Expression patterns of calretinin, calbindin and parvalbumin and their colocalization in neurons during development of Macaca monkey retina.
Hendrickson A et al. Exp. Eye Res. (2007) PubMed:17845803

Ultrastructural study of gap junctions between dendrites of parvalbumin-containing GABAergic neurons in various neocortical areas of the adult rat.
Fukuda T et al. Neuroscience (2003) PubMed:12849736

Calcium-binding protein parvalbumin-immunoreactive neurons in the rat olfactory bulb. 2. Postnatal development.
Kosaka K et al. Exp Brain Res (1994) PubMed:7925803

Immunocytochemical localization of the plasma membrane calcium pump, calbindin-D28k, and parvalbumin in Purkinje cells of avian and mammalian cerebellum.
Tolosa de Talamoni N et al. Proc. Natl. Acad. Sci. U.S.A. (1993) PubMed:8265654

Neostriatal GABAergic interneurons contain NOS, calretinin or parvalbumin.
Kubota Y et al. Neuroreport (1993) PubMed:7507722

Access the online factsheet including applicable protocols at <https://sysy.com/product/195308> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.