

VAMP7

Cat.No. 232 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin and azide were added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP staining) IP: not recommended ICC: not recommended IHC: 1 : 500 IHC-P: not tested yet
Immunogen	Recombinant protein corresponding to AA 10 to 177 from mouse VAMP7 (UniProt Id: P70280)
Reactivity	Reacts with: rat (Q9JHW5), mouse (P70280), human (P51809). No signal: zebrafish. Other species not tested yet.
Specificity	K.O. validated PubMed: 30815249
Matching control	232-0P

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

VAMP 7, also referred to as Ti-VAMP and SybL 1, is a member of the SNARE family of proteins and a relative of synaptobrevin. It is involved in membrane fusion events that mediate neurite outgrowth in developing neurons, in endosome to lysosome transport and in other cellular trafficking mechanisms. VAMP 7 is ubiquitously expressed in different tissues. It is a member of the syntaxin 4-SNAP 23-VAMP 7- and the syntaxin 7-syntaxin 8-Vti1b-VAMP 7-SNARE complex.

Selected References for 232 003

- Comparative study of commercially available and homemade anti-VAMP7 antibodies using CRISPR/Cas9-depleted HeLa cells and VAMP7 knockout mice.
Verraes A, Cholley B, Galli T, Nola S
F1000Research (2018) 7: 1649. . **WB; KO verified; tested species: human**
- Proteomic analysis reveals the composition of glutamatergic organelles of auditory inner hair cell.
Cepeda AP, Ninov M, Neef J, Parfentev I, Kusch K, Reisinger E, Jahn R, Moser T, Urlaub H
Molecular & cellular proteomics : MCP (2023) : 100704. . **IHC; tested species: mouse**
- Single synapse glutamate imaging reveals multiple levels of release mode regulation in mammalian synapses.
Farsi Z, Walde M, Klementowicz AE, Paraskevopoulou F, Woehler A
iScience (2021) 241: 101909. . **ICC; tested species: rat**
- Syntaxin 11 binds Vti1b and regulates late endosome to lysosome fusion in macrophages.
Offenhäuser C, Lei N, Roy S, Collins BM, Stow JL, Murray RZ
Traffic (Copenhagen, Denmark) (2011) 126: 762-73. . **WB**
- Lysosomal exocytosis releases pathogenic α -synuclein species from neurons in synucleinopathy models.
Xie YX, Naseri NN, Fels J, Kharel P, Na Y, Lane D, Burré J, Sharma M
Nature communications (2022) 131: 4918. . **WB; tested species: mouse**
- Characterisation of GLUT4 trafficking in HeLa cells: comparable kinetics and orthologous trafficking mechanisms to 3T3-L1 adipocytes.
Morris S, Geoghegan ND, Sadler JBA, Koester AM, Black HL, Laub M, Miller L, Heffernan L, Simpson JC, Mastick CC, Cooper J, et al.
PeerJ (2020) 8: e8751. . **WB; tested species: mouse**
- VAMP4 maintains a Ca²⁺-sensitive pool of spontaneously recycling synaptic vesicles.
Lin PY, Chanaday NL, Horvath PM, Ramirez DMO, Monteggia LM, Kavalali ET
The Journal of neuroscience : the official journal of the Society for Neuroscience (2020) : . . **WB; tested species: rat**
- A trap mutant reveals the physiological client spectrum of TRC40.
Coy-Vergara J, Rivera-Monroy J, Urlaub H, Lenz C, Schwappach B
Journal of cell science (2019) 13213: . . **WB; tested species: human**
- Cardiac SNARE Expression in Health and Disease.
Bowman PRT, Smith GL, Gould GW
Frontiers in endocrinology (2019) 10: 881. . **WB; tested species: mouse**
- Selective molecular impairment of spontaneous neurotransmission modulates synaptic efficacy.
Crawford DC, Ramirez DM, Trauterman B, Monteggia LM, Kavalali ET
Nature communications (2017) 8: 14436. . **WB; KD verified**
- Differential Expression of Munc13-2 Produces Unique Synaptic Phenotypes in the Basolateral Amygdala of C57BL/6J and DBA/2J Mice.
Gioia DA, Alexander NJ, McCool BA
The Journal of neuroscience : the official journal of the Society for Neuroscience (2016) 3643: 10964-10977. . **WB**

Access the online factsheet including applicable protocols at <https://sysy.com/product/232003> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.