

SynCAM2

Cat.No. 243 203; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin was added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP staining) IP: not tested yet ICC: 1 : 100 IHC: 1 : 100 IHC-P: not tested yet
Immunogen	Synthetic peptide corresponding to AA 167 to 181 from mouse SynCAM2 (UniProt Id: Q8BLQ9)
Reactivity	Reacts with: rat (Q1WIM2), mouse (Q8BLQ9), human (Q8N3J6). Other species not tested yet.
Specificity	K.O. validated PubMed: 29217450
Matching control	243-2P

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Development of synapses involves transsynaptic interactions of dedicated synaptic adhesion molecules like neuroligins, neurexins and synCAMs (**Synaptic cell adhesion molecules**). Four SynCAM isoforms have been described so far. All share a common domain structure and contain three extracellular Ig domains, a single transmembrane region, and a short COOH-terminal cytoplasmic tail. **SynCAM 2** has been shown to be associated to myelinated axons and **synCAM 3** can serve as a receptor for herpes viruses.

Selected References for 243 203

Cadm2 regulates body weight and energy homeostasis in mice.
Yan X, Wang Z, Schmidt V, Gauer A, Willnow TE, Heinig M, Poy MN
Molecular metabolism (2018) 8: 180-188. . **WB, ICC; KO verified; tested species: mouse**

Regulation of body weight and energy homeostasis by neuronal cell adhesion molecule 1.
Rathjen T, Yan X, Kononenko NL, Ku MC, Song K, Ferrarese L, Tarallo V, Puchkov D, Kochlamazashvili G, Brachs S, Varela L, et al.
Nature neuroscience (2017) 208: 1096-1103. . **ICC; KO verified; tested species: mouse**

Selected General References

Aberrations of a cell adhesion molecule CADM4 in renal clear cell carcinoma.
Nagata M, Sakurai-Yageta M, Yamada D, Goto A, Ito A, Fukuhara H, Kume H, Morikawa T, Fukayama M, Homma Y, Murakami Y, et al.
International journal of cancer (2012) 1306: 1329-37. .

Localization of Cadm2a and Cadm3 proteins during development of the zebrafish nervous system.
Hunter PR, Nikolaou N, Odermatt B, Williams PR, Drescher U, Meyer MP
The Journal of comparative neurology (2011) 51911: 2252-70. .

The cell adhesion nectin-like molecules (Nectin) 1 and 4 suppress the growth and tumorigenic ability of colon cancer cells.
Raveh S, Gavert N, Spiegel I, Ben-Ze'ev A
Journal of cellular biochemistry (2009) 1081: 326-36. .

The adhesion molecule Nectin-3/SynCAM-2 localizes to myelinated axons, binds to oligodendrocytes and promotes cell adhesion.
Pellissier F, Gerber A, Bauer C, Ballivet M, Ossipow V
BMC neuroscience (2007) 8: 90. .

Nectin-like proteins mediate axon Schwann cell interactions along the internode and are essential for myelination.
Maurel P, Einheber S, Galinska J, Thaker P, Lam I, Rubin MB, Scherer SS, Murakami Y, Gutmann DH, Salzer JL
The Journal of cell biology (2007) 1785: 861-74. .

SynCAMs organize synapses through heterophilic adhesion.
Fogel AI, Akins MR, Krupp AJ, Stagi M, Stein V, Biederer T
The Journal of neuroscience : the official journal of the Society for Neuroscience (2007) 2746: 12516-30. .

Crystal structure of the V domain of human Nectin-like molecule-1/Syncam3/Tsll1/Igsf4b, a neural tissue-specific immunoglobulin-like cell-cell adhesion molecule.
Dong X, Xu F, Gong Y, Gao J, Lin P, Chen T, Peng Y, Qiang B, Yuan J, Peng X, Rao Z, et al.
The Journal of biological chemistry (2006) 28115: 10610-7. .

Nectin-like molecule-1/TSLL1/SynCAM3: a neural tissue-specific immunoglobulin-like cell-cell adhesion molecule localizing at non-junctional contact sites of presynaptic nerve terminals, axons and glia cell processes.
Kakunaga S, Ikeda W, Itoh S, Deguchi-Tawarada M, Ohtsuka T, Mizoguchi A, Takai Y
Journal of cell science (2005) 118Pt 6: 1267-77. .

SynCAM, a synaptic adhesion molecule that drives synapse assembly.
Biederer T, Sara Y, Mozhayeva M, Atasoy D, Liu X, Kavalali ET, Südhof TC
Science (New York, N.Y.) (2002) 2975586: 1525-31. .

Access the online factsheet including applicable protocols at <https://sysy.com/product/243203> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.