

GAPDH

Cat.No. 247 002; Polyclonal rabbit antibody, 200 µl antiserum (lyophilized)

Data Sheet

Reconstitution/ Storage	200 µl antiserum, lyophilized. For reconstitution add 200 µl H ₂ O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 up to 1 : 5000 (AP staining) IP: not tested yet ICC: not tested yet IHC: not tested yet IHC-P: not tested yet Clarity: external data (see remarks)
Immunogen	Recombinant protein corresponding to AA 1 to 335 from human GAPDH (UniProt Id: P04406)
Reactivity	Reacts with: human (P04406), rat (P04797), mouse (P16858), zebrafish. Other species not tested yet.
Remarks	Well established as a loading control for Western-blotting. Clarity: This antibody has been successfully used for CLARITY application in human brain (Woelfle et al., 2023; PMID: 37221592).

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Glyceraldehyde 3-phosphate dehydrogenase (GAPDH), also referred to as G3PDH, catalyzes the sixth step of glycolysis and thus plays a crucial role in the glucose metabolism. In addition to this well known metabolic function it also exhibits nitrosylase activities and is involved in cellular events like transcription, RNA transport, DNA replication and apoptosis. It is an abundant housekeeping gene and is frequently used as a loading reference in westernblot experiments.

Selected References for 247 002

Interactome of the amyloid precursor protein APP in brain reveals a protein network involved in synaptic vesicle turnover and a close association with Synaptotagmin-1.

Kohli BM, Pflieger D, Mueller LN, Carbonetti G, Aebersold R, Nitsch RM, Konietzko U
Journal of proteome research (2012) 118: 4075-90. . **WB**

Computerized EEG in the comparison of oxyprothepin and fluphenazine decanoate.

Misurec J, Náhunek K, Svestka J, Cesková E

Activitas nervosa superior (1979) 213: 140. . **CLARITY; tested species: human**

PHF2-mediated H3K9me balance orchestrates heterochromatin stability and neural progenitor proliferation.

Aguirre S, Pappa S, Serna-Pujol N, Padilla N, Iacobucci S, Nacht AS, Vicent GP, Jordan A, de la Cruz X, Martínez-Balbás MA
EMBO reports (2024) : . . **WB; tested species: mouse**

Oleoylethanolamide and Palmitoylethanolamide Enhance IFNβ-Induced Apoptosis in Human Neuroblastoma SH-SY5Y Cells.

Camoglio C, Balla J, Fadda P, Dedoni S

Molecules (Basel, Switzerland) (2024) 297: . . **WB; tested species: human**

Loss of interleukin 1 signaling causes impairment of microglia- mediated synapse elimination and autistic-like behaviour in mice.

Borrecia A, Mantovani C, Desiato G, Corradini I, Filipello F, Elia CA, D'Autilia F, Santamaria G, Garlanda C, Morini R, Pozzi D, et al.
Brain, behavior, and immunity (2024) 117: 493-509. . **WB; tested species: mouse**

Changes in glial cell activation and extracellular vesicles production precede the onset of disease symptoms in transgenic hSOD1G93A pigs.

Golia MT, Frigerio R, Pucci S, Sironi F, Margotta C, Pasetto L, Testori C, Berrone E, Ingravalle F, Chiari M, Gori A, et al.

Experimental neurology (2024) 374: 114716. . **WB; tested species: pig**

Lysophosphatidic Acid Stimulates Mitogenic Activity and Signaling in Human Neuroblastoma Cells through a Crosstalk with Anaplastic Lymphoma Kinase.

Dedoni S, Olinas MC, Onali P

Biomolecules (2024) 146: . . **WB; tested species: human**

Microglial large extracellular vesicles propagate early synaptic dysfunction in Alzheimer's disease.

Gabrielli M, Prada I, Joshi P, Falcicchia C, D'Arrigo G, Rutigliano G, Battocchio E, Zenatelli R, Tozzi F, Radeghieri A, Arancio O, et al.

Brain : a journal of neurology (2022) : . . **WB; tested species: mouse**

Prenatal interleukin 6 elevation increases glutamatergic synapse density and disrupts hippocampal connectivity in offspring.

Mirabella F, Desiato G, Mancinelli S, Fossati G, Rasile M, Morini R, Markicevic M, Grimm C, Amegandjin C, Termanini A, Peano C, et al.

Immunity (2021) 5411: 2611-2631.e8. . **WB; tested species: mouse**

The histone demethylase PHF8 regulates astrocyte differentiation and function.

Iacobucci S, Padilla N, Gabrielli M, Navarro C, Lombardi M, Vicioso-Mantis M, Verderio C, de la Cruz X, Martínez-Balbás MA
Development (Cambridge, England) (2021) 14812: . . **WB; tested species: mouse**

Nrg1 haploinsufficiency alters inhibitory cortical circuits.

Navarro-Gonzalez C, Carceller H, Benito Vicente M, Serra I, Navarrete M, Domínguez-Canterla Y, Rodríguez-Prieto Á, González-Manteiga A, Fazzari P

Neurobiology of disease (2021) 157: 105442. . **WB; tested species: mouse**

Access the online factsheet including applicable protocols at <https://sysy.com/product/247002> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.