

Giantin

Cat.No. 263 005; Polyclonal Guinea pig antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin and azide were added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP staining) IP: not tested yet ICC: 1 : 500 IHC: 1 : 200 up to 1 : 500 IHC-P: 1 : 200 up to 1 : 500
Immunogen	Recombinant protein corresponding to AA 1 to 427 from rat Giantin (UniProt Id: G3V6A8)
Reactivity	Reacts with: rat (G3V6A8), mouse. Other species not tested yet.
Remarks	WB: Due to the large size of this protein, we recommend NuPAGE 3-8% Tris-Acetate gels for SDS-PAGE.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

In eukaryotic cells the Golgi complex consists of stacked cisternae. **Giantin**, also referred to as **macroglglin** or **p400**, is a single pass membrane protein involved in tethering these cisternae by cross-bridge formation. Its large cytoplasmic N-terminus is involved in the docking of COPI vesicles via p115 to the Golgi membrane.

Systemic lupus erythematosus and Sjögren's syndrome correlate with autoimmune antibodies against Golgi-compartment proteins with giantin being the most common autoantigen.

Selected References for 263 005

A trap mutant reveals the physiological client spectrum of TRC40.
Coy-Vergara J, Rivera-Monroy J, Urlaub H, Lenz C, Schwappach B
Journal of cell science (2019) 13213: . . **ICC; tested species: human**

The X-Linked Intellectual Disability Gene Zdhc9 Is Essential for Dendrite Outgrowth and Inhibitory Synapse Formation.
Shimell JJ, Shah BS, Cain SM, Thouta S, Kuhlmann N, Tatarnikov I, Jovellar DB, Brigidi GS, Kass J, Milnerwood AJ, Snutch TP, et al.
Cell reports (2019) 298: 2422-2437.e8. . **ICC; tested species: rat**

Selected General References

Giantin interacts with both the small GTPase Rab6 and Rab1.
Rosing M et al. Exp. Cell Res. (2007) PubMed:17475246

Giantin is the major Golgi autoantigen in human anti-Golgi complex sera.
Nozawa K et al. Arthritis Res. Ther. (2004) PubMed:15059272

The p115-interactive proteins GM130 and giantin participate in endoplasmic reticulum-Golgi traffic.
Alvarez C et al. J. Biol. Chem. (2001) PubMed:11035033

A role for giantin in docking COPI vesicles to Golgi membranes.
Sönnichsen B et al. J. Cell Biol. (1998) PubMed:9490716

Molecular genetic analyses of a 376-kilodalton Golgi complex membrane protein (giantin)
Seelig HP et al. Mol. Cell. Biol. (1994) PubMed:7511208

Giantin, a novel conserved Golgi membrane protein containing a cytoplasmic domain of at least 350 kDa.
Linstedt AD et al. Mol. Biol. Cell (1993) PubMed:7691276

Access the online factsheet including applicable protocols at <https://sysy.com/product/263005> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.