

WRB

Cat.No. 324 103; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin was added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP staining) IP: yes ICC: not tested yet IHC: not tested yet IHC-P: not tested yet
Immunogen	Recombinant protein corresponding to AA 30 to 99 from human WRB (UniProt Id: O00258)
Reactivity	Reacts with: rat (Q6P6S5), mouse (Q8K0D7), human (O00258). Other species not tested yet.
Specificity	K.D. validated PubMed: 27226539

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

The widely expressed tryptophan (**W**)-rich basic protein (**WRB**), also known as congenital heart disease protein 5 (**CHD 5**) has initially been described as a nuclear protein that plays a potential role in the pathogenesis of CHD in down syndrome patients. Recently this protein has been identified as the receptor for TRC40/Asna1-mediated insertion of tail-anchored proteins into the ER membrane.

Selected References for 324 103

The WRB Subunit of the Get3 Receptor is Required for the Correct Integration of its Partner CAML into the ER. Carvalho HJF, Del Bondio A, Maltecca F, Colombo SF, Borgese N. Scientific reports (2019) 91: 11887. . **WB, IP; tested species: human**

ER entry pathway and glycosylation of GPI-anchored proteins are determined by N-terminal signal sequence and C-terminal GPI-attachment sequence.

Hirata T, Yang J, Tomida S, Tokoro Y, Kinoshita T, Fujita M, Kizuka Y. The Journal of biological chemistry (2022) : 102444. . **WB; tested species: human**

Discrimination between the endoplasmic reticulum and mitochondria by spontaneously inserting tail-anchored proteins. Figueiredo Costa B, Cassella P, Colombo SF, Borgese N. Traffic (Copenhagen, Denmark) (2018) 193: 182-197. . **WB**

Tail-anchored Protein Insertion in Mammals: FUNCTION AND RECIPROCAL INTERACTIONS OF THE TWO SUBUNITS OF THE TRC40 RECEPTOR.

Colombo SF, Cardani S, Maroli A, Vitiello A, Soffientini P, Crespi A, Bram RF, Benfante R, Borgese N. The Journal of biological chemistry (2016) 29129: 15292-306. . **WB; KD verified; tested species: rat**

Selected General References

WRB and CAML are necessary and sufficient to mediate tail-anchored protein targeting to the ER membrane.

Vilardi F, Stephan M, Clancy A, Janshoff A, Schwappach B. PloS one (2014) 91: e85033. .

WRB is the receptor for TRC40/Asna1-mediated insertion of tail-anchored proteins into the ER membrane.

Vilardi F, Lorenz H, Dobberstein B. Journal of cell science (2011) 124Pt 8: 1301-7. .

Identification and characterization of a new human cDNA from chromosome 21q22.3 encoding a basic nuclear protein. Egeo A, Mazzocco M, Sotgia F, Arrigo P, Oliva R, Bergonon S, Nizetic D, Rasore-Quartino A, Scartezzini P. Human genetics (1998) 1023: 289-93. .

Access the online factsheet including applicable protocols at <https://susy.com/product/324103> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.