

## Somatostatin-28

Cat.No. 366 004; Polyclonal Guinea pig antibody, 100 µl antiserum (lyophilized)

### Data Sheet

|                            |   |
|----------------------------|---|
| Reconstitution/<br>Storage | 100 µl antiserum, lyophilized. For <b>reconstitution</b> add 100 µl H <sub>2</sub> O, then aliquot and store at -20°C until use.<br>Antibodies should be stored at +4°C when still lyophilized. Do not freeze!<br>For detailed information, see back of the data sheet. |
| Applications               | <b>WB:</b> not tested yet<br><b>IP:</b> not tested yet<br><b>ICC:</b> 1 : 500<br><b>IHC:</b> 1 : 500<br><b>IHC-P:</b> 1 : 500   |
| Immunogen                  | Synthetic peptide corresponding to AA 89 to 100 from mouse Somatostatin (UniProt Id: P60041)  |
| Reactivity                 | Reacts with: human (P61278), rat (P60042), mouse (P60041).<br>Other species not tested yet.   |
| Specificity                | This antibody preferentially recognizes somatostatin-28. It only shows minor cross-reactivity to the unprocessed precursor protein and does not detect somatostatin-14.   |

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

### Background

**Somatostatin**, also referred to as **SST**, **growth hormone-inhibiting hormone** or **GHIH**, is a peptide hormone that regulates the endocrine system and affects neurotransmission and cell proliferation via interaction with G protein-coupled somatostatin receptors. It inhibits the secretion of many important hormones, including insulin, glucagon and somatotropin (also designated growth hormone, or GH). Somatostatin has two forms, active 14 amino acid and 28 amino acid. They are produced by alternative cleavage of the single precursor protein encoded by this gene.

### Selected References for 366 004

Generation of self-organized autonomic ganglion organoids from fibroblasts.

Liu S, Xiang K, Yuan F, Xiang M

iScience (2023) 263: 106241. . **ICC; tested species: mouse**

Amyloid β induces interneuron-specific changes in the hippocampus of APPNL-F mice.

Sos KE, Mayer MI, Takács VT, Major A, Bardóczi Z, Beres BM, Szeles T, Saito T, Saido TC, Mody I, Freund TF, et al.

PLoS one (2020) 155: e0233700. . **EM; tested species: mouse**

Phase-specific surround suppression in mouse primary visual cortex correlates with figure detection behavior based on phase discontinuity.

Li F, Jiang W, Wang TY, Xie T, Yao H

Neuroscience (2018) : . . **IHC; tested species: mouse**

Synaptic and dendritic architecture of different types of hippocampal somatostatin interneurons.

Takács V, Bardóczi Z, Orosz Á, Major A, Tar L, Berki P, Papp P, Mayer MI, Sebők H, Zsolt L, Sos KE, et al.

PLoS biology (2024) 223: e3002539. . **IHC; tested species: mouse**

Selective prosaposin expression in Langerhans islets of the mouse pancreas.

Fuyuki A, Sohel MSH, Homma T, Kitamura K, Takashima S, Onouchi S, Saito S

Tissue & cell (2024) 88: 102367. . **IHC; tested species: mouse**

Voluntary running-induced activation of ventral hippocampal GABAergic interneurons contributes to exercise-induced hypoalgesia in neuropathic pain model mice.

Minami K, Kami K, Nishimura Y, Kawanishi M, Imashiro K, Kami T, Habata S, Senba E, Umemoto Y, Tajima F

Scientific reports (2023) 131: 2645. . **IHC; tested species: mouse**

PDGF-BB-Dependent Neurogenesis Buffers Depressive-Like Behaviors by Inhibition of GABAergic Projection from Medial Septum to Dentate Gyrus.

Li HH, Liu Y, Chen HS, Wang J, Li YK, Zhao Y, Sun R, He JG, Wang F, Chen JG

Advanced science (Weinheim, Baden-Württemberg, Germany) (2023) 1022: e2301110. . **IHC; tested species: mouse**

Nrg1 haploinsufficiency alters inhibitory cortical circuits.

Navarro-Gonzalez C, Carceller H, Benito Vicente M, Serra I, Navarrete M, Domínguez-Canterla Y, Rodríguez-Prieto Á, González-Manteiga A, Fazzari P

Neurobiology of disease (2021) 157: 105442. . **IHC; tested species: mouse**

Total Number and Ratio of GABAergic Neuron Types in the Mouse Lateral and Basal Amygdala.

Vereczki VK, Müller K, Krizsán É, Máté Z, Fekete Z, Rovira-Esteban L, Veres JM, Erdélyi F, Hájos N

The Journal of neuroscience : the official journal of the Society for Neuroscience (2021) 4121: 4575-4595. . **IHC; tested species: mouse**

ROCK/PKA inhibition rescues hippocampal hyperexcitability and GABAergic neuron alterations in Oligophrenin-1 Knock-out mouse model of X-linked intellectual disability.

Busti I, Allegra M, Spalletti C, Panzi C, Restani L, Billuart P, Caleo M

The Journal of neuroscience : the official journal of the Society for Neuroscience (2020) : . . **IHC; tested species: mouse**

Glia-to-Neuron Conversion by CRISPR-CasRx Alleviates Symptoms of Neurological Disease in Mice.

Zhou H, Su J, Hu X, Zhou C, Li H, Chen Z, Xiao Q, Wang B, Wu W, Sun Y, Zhou Y, et al.

Cell (2020) : . . **IHC; tested species: mouse**

Access the online factsheet including applicable protocols at <https://sysy.com/product/366004> or scan the QR-code.



## FAQ - How should I store my antibody?

### Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

### Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

### Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

### Product Specific Hints for Storage

#### Control proteins / peptides

- Store at -20°C to -80°C.

#### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

## Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.