

Dopamine receptor D2

Cat.No. 376 203; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin and azide were added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP staining) IP: not tested yet ICC: 1 : 500 IHC: 1 : 500 IHC-P: 1 : 500
Immunogen	Recombinant protein corresponding to AA 222 to 368 from mouse DRD2 (UniProt Id: P61168)
Reactivity	Reacts with: rat (P61169), mouse (P61168). Other species not tested yet.
Specificity	K.O. validated PubMed: 36170827

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Dopamine receptors transduce the signal of the neurotransmitter dopamine. Dopamine regulates a variety of functions including locomotor activity, emotion, food intake, hormone secretion, learning and memory. The dysregulation of the dopaminergic system results in several neurological and neuropsychiatric diseases including Parkinson's disease, dystonia, and schizophrenia. Dopamine plays also an important role in the pathogenesis of hypertension by regulating epithelial sodium transport and by interacting with vasoactive hormones.

All five dopamine receptors belong to the 7-transmembrane domain, G protein-coupled receptor superfamily. They have been divided into two subfamilies: Two D1-like receptor subtypes (**D1** and **D5**) couple to the G protein Gs and activate adenylyl cyclase, increasing the intracellular concentration of the second messenger cAMP. The other receptor subtypes belong to the D2-like subfamily (**D2**, **D3**, and **D4**) and are prototypic of G protein-coupled receptors that inhibit adenylyl cyclase.

Dopamine receptors can form heteromeric complexes with dopamine receptors from other subtypes or with receptors of other endogenous signaling ligands. These heteromeric complexes have functional properties distinct from the component receptors or are able to modulate the canonical signaling.

Dopamine receptor D1 is widely distributed throughout the brain with the highest expression in the striatum. In the periphery, the D1 receptor has been detected in the adrenal cortex, kidney and heart. Recently, it was shown that dopamine receptor D1 is expressed in breast cancer, thereby identifying this receptor as a novel therapeutic target in this disease. D1 receptor overexpression is associated with advanced breast cancer and poor prognosis.

Dopamine Receptor D2 is most abundant in the striatum where it is expressed in medium spiny neurons. Functionally, the D1 and D2 Receptors have been implicated in the regulation of both locomotor and cognitive functions.

Selected References for 376 203

- Nanosopic dopamine transporter distribution and conformation are inversely regulated by excitatory drive and D2 autoreceptor activity.
Lycas MD, Ejdrup AL, Sørensen AT, Haahr NO, Jørgensen SH, Guthrie DA, Støier JF, Werner C, Newman AH, Sauer M, Herborg F, et al.
Cell reports (2022) 4013: 111431. . **ICC, IHC; KO verified; tested species: mouse, rat**
- Loss of the parkinsonism-associated protein FBXO7 in glutamatergic forebrain neurons in mice leads to abnormal motor behavior and synaptic defects.
Wang J, Joseph S, Vingill S, Dere E, Tatenhorst L, Ronnenberg A, Lingor P, Preisinger C, Ehrenreich H, Schulz JB, Stegmüller J, et al.
Journal of neurochemistry (2023) : . . **IHC; tested species: mouse**

Selected General References

- Expression and therapeutic targeting of dopamine receptor-1 (D1R) in breast cancer.
Borcherding DC, Tong W, Hugo ER, Barnard DF, Fox S, LaSance K, Shaughnessy E, Ben-Jonathan N
Oncogene (2016) 3524: 3103-13. .
- Dopamine receptor heteromeric complexes and their emerging functions.
George SR, Kern A, Smith RG, Franco R
Progress in brain research (2014) 211: 183-200. .
- Dopamine receptor mapping with PET imaging in Parkinson's disease.
Niccolini F, Su P, Politis M
Journal of neurology (2014) 26112: 2251-63. .

Access the online factsheet including applicable protocols at <https://sysy.com/product/376203> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.