

Galanin

Cat.No. 446 004; Polyclonal Guinea pig antibody, 100 µl antiserum (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µl antiserum, lyophilized. For reconstitution add 100 µl H ₂ O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: not tested yet IP: not tested yet ICC: 1 : 500 IHC: 1 : 500 IHC-P: 1 : 500 up to 1 : 2500
Immunogen	Synthetic peptide corresponding to AA 33 to 61 from mouse Galanin (UniProt Id: P47212)
Reactivity	Reacts with: mouse (P47212), rat (P10683). Other species not tested yet.
Specificity	The antibody is specific for galanin. It may crossreact with the unprocessed precursor protein and with galanin-like peptide due to sequence homology.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Galanin, also referred to as Gal or Galn, is a 30 amino acid neuropeptide (29 in rodents) which was first isolated from porcine intestine in 1983 (1). It is synthesized as part of a larger propeptide which is proteolytically processed to release the mature neuropeptide. Galanin is highly conserved across species and has a widespread presence in the periphery and the central nervous system, including brain, spinal cord, and gastrointestinal tract. The effects of this neuropeptide are mediated through three G protein coupled galanin receptors (2, 3).

Galanin has been implicated in many biologically diverse functions including arousal/sleep regulation, nociception, cognition, feeding, reproduction, osmotic homeostasis, glucose metabolism, and regulation of mood (3, 4).

Several human diseases, including Alzheimer's disease, epilepsy, diabetes mellitus and mood disorders like anxiety and depression are associated with the disturbance of the galaninergic system (2, 3).

Selected General References

Galanin - a novel biologically active peptide from porcine intestine.
Tatemoto K, Rökæus A, Jörnvall H, McDonald TJ, Mutt V
FEBS letters (1983) 1641: 124-8. .

Neuropeptide modulation of addiction: Focus on galanin.
Genders SG, Scheller KJ, Djouma E
Neuroscience and biobehavioral reviews (2020) 110: 133-149. .

Galanin peptide family regulation of glucose metabolism.
Fang P, Yu M, Shi M, Bo P, Zhang Z
Frontiers in neuroendocrinology (2020) 56: 100801. .

Physiology, signaling, and pharmacology of galanin peptides and receptors: three decades of emerging diversity.
Lang R, Gundlach AL, Holmes FE, Hobson SA, Wynick D, Hökfelt T, Kofler B
Pharmacological reviews (2015) 67: 118-75. .

Access the online factsheet including applicable protocols at <https://sysy.com/product/446004> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.