

TMEM106B

Cat.No. 506 017; Monoclonal rat antibody, 100 µg purified IgG (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µg purified IgG, lyophilized. Albumin and azide were added for stabilization. For reconstitution add 100 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (see remarks) IP: not tested yet ICC: not tested yet IHC: not tested yet IHC-P: 1 : 1000 up to 1 : 2000 (see remarks)
Clone	SY-118C4
Subtype	IgG2a (κ light chain)
Immunogen	Synthetic peptide corresponding to AA 239 to 252 from human TMEM106B (UniProt Id: Q9NUM4)
Epitop	AA 239 to 252 from human TMEM106B (UniProt Id: Q9NUM4)
Reactivity	Reacts with: human (Q9NUM4), mouse (Q80X71). Other species not tested yet.
Specificity	K.O. validated
Remarks	WB: This antibody shows an additional unspecific band at 70kD. IHC-P: Antigen retrieval with Tris-EDTA buffer pH 9, followed by formic acid treatment, is required. Treat with 80% formic acid for 10 min

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

TMEM106B or Transmembrane protein 106B is a lysosomal type II transmembrane protein which has been identified by a genome-wide association study as a potential risk factor for a neurodegenerative disorder called frontotemporal lobar degeneration (FTLD)(1). The C-terminal luminal domain of TMEM106B undergoes a proteolytic cleavage that forms sarkosyl-insoluble amyloid-like fibrils (residues 120-254) in human brains (2). TMEM106B is ubiquitously expressed, but the function of TMEM106B is almost unknown. TMEM106B is a receptor mediating ACE2-independent SARS-CoV-2 cell entry (3).

Selected References for 506 017

Physiological shedding and C-terminal proteolytic processing of TMEM106B.
Held S, Erck C, Kemppainen S, Bleibaum F, Giridhar NJ, Feederle R, Krenner C, Juopperi SP, Calliari A, Mentrup T, Schröder B, et al.
Cell reports (2024) 441: 115107. . **WB, IHC-P; tested species: human,mouse**

Selected General References

Common variants at 7p21 are associated with frontotemporal lobar degeneration with TDP-43 inclusions.
Van Deerlin VM et al. Nat Genet (2010) PubMed:20154673
TMEM106B is a receptor mediating ACE2-independent SARS-CoV-2 cell entry.
Baggen J et al. Cell (2023) PubMed:37421949
Age-dependent formation of TMEM106B amyloid filaments in human brains.
Schweighauser M et al. Nature (2022) PubMed:35344985

Access the online factsheet including applicable protocols at <https://sysy.com/product/506017> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.